

## Research Note

# An Anomaly (Gynandromorphism) in *Abbreviata* sp. (Nematoda: Physalopteridae)

JOHN H. GREVE

Iowa State University, Ames, Iowa 50011 (e-mail: mehull@iastate.edu)

**ABSTRACT:** A single gynandromorphous *Abbreviata* sp. was recovered from the stomach of a black tree monitor, *Varanus beccarii*, from the Aru Islands, Indonesia.

**KEY WORDS:** gynandromorph, *Varanus*, anomaly, hermaphrodite, *Abbreviata*.

Anomalies in nematodes are seldom seen. Among anomalies reported for nematodes, those occurring in the reproductive system are by far the most common (Belogurov and Belogurova, 1979). In a review of 14 reports (1924–1989) on anomalies in nematodes, 10 dealt with abnormalities of the reproductive system (spicules, gubernacula, bursal rays, uteri, ovaries, and eggs) (Becklund, 1960; Andrews, 1970; Goldstein, 1977; Amin, 1989), whereas the other 4 dealt with abnormalities of the musculature or the digestive system (Lyons and Goldsmid, 1973); however, none described a gynandromorph. Most reported anomalies involve ascarids or strongylates, which possibly reflects frequency with which these nematodes are examined relative to other kinds of nematodes.

A black tree monitor (*Varanus beccarii*) had been wild-caught in the Aru Islands, Indonesia, and illegally shipped to the United States. When confiscated by authorities, the monitor was in very poor physical condition, and it died shortly thereafter. At necropsy, a single nematode specimen was found in the stomach and submitted to the Department of Veterinary Pathology, Iowa State University, for identification. After clearing in Hoyer's solution (Baker et al., 1956), the nematode was discovered to be a gynandromorphous *Abbreviata* sp. (Fig. 1). It was assumed that this bisexual condition represented an anomaly, because gynandromorphism is not a characteristic of nematodes and no gynandromorphs were mentioned in specimens of *Abbreviata* spp. commonly found in monitors (Jones, 1988).

The gynandromorph had a single uterus with a long vagina opening near the cloaca (precise site of this opening could not be seen). The uterus and vagina contained small (44 by 25  $\mu$ m) embryonated eggs typical for the genus. Male com-

ponents included a typical physalopterine bursa and dissimilar, unequal spicules (right = 350  $\mu$ m, left = 2,200  $\mu$ m). The parasite has been deposited in the U.S. National Parasite Collection (Accession No. 85054).




Figure 1. Outline drawing of posterior portion of gynandromorphous *Abbreviata* sp. from black tree monitor. Highlighted features are uterus (A), left spicule (B), and right spicule (C). Scale bar = 500  $\mu$ m.

Furthermore, the finding of *Abbreviata* sp. in *Varanus beccarii* appears to be a new host record (Baker, 1987).

The author thanks Thomas P. Alvarado, D.V.M., Dallas, Texas for submitting this specimen.

### Literature Cited

- Amin, O. M. 1989. Abnormalities in some helminth parasites of fish. Transactions of the American Microscopical Society 108:27-39.
- Andrews, J. R. H. 1970. The teratology of spicules in some trichostrongyline nematodes (family Trichostrongylidae Leiper, 1912). Journal of Parasitology 56:402-404.
- Baker, E. W., T. M. Evans, D. J. Gould, W. B. Hull, and H. L. Keegan. 1956. A Manual of Parasitic Mites of Medical or Economic Importance. Technical Publication, National Pest Control Association, New York. 170 pp.
- Baker, M. R. 1987. Synopsis of the Nematoda Parasitic in Amphibians and Reptiles. Memorial University of Newfoundland Occasional Papers in Biology, No. 11. 325 pp.
- Becklund, W. W. 1960. Morphologic anomalies in male *Haemonchus contortus* (Rudolphi, 1803) Cobb, 1898 (Nematoda: Trichostrongylidae) from sheep. Proceedings of the Helminthological Society of Washington 27:194-199.
- Belogurov, O. I., and L. S. Belogurova. 1979. On morphological abnormalities (deformities) in nematodes and their possible evolutionary value. Zoologicheskii Zhurnal 58:1730-1733 (in Russian).
- Goldstein, P. 1977. Incidence of an extra reproductive tract in female *Ascaris suum*. Journal of Parasitology 63:689.
- Jones, H. I. 1988. Nematodes from nine species of *Varanus* (Reptilia) from tropical northern Australia, with particular reference to the genus *Abbreviata* (Phyllopteridae). Australian Journal of Zoology 36:691-708.
- Lyons, N. F., and J. M. Goldsmid. 1973. Abnormal *Ternidens deminutus* Railliet and Henry, 1909 (Nematoda) from man in Rhodesia. Journal of Parasitology 59:219-220.

J. Helminthol. Soc. Wash.  
63(2), 1996, pp. 262-265

### Research Note

## Helminth Parasites of the Osprey, *Pandion haliaetus*, in North America

JOHN M. KINSELLA,<sup>1,3</sup> REBECCA A. COLE,<sup>2</sup> DONALD J. FORRESTER,<sup>1</sup>  
AND C. L. RODERICK<sup>2</sup>

<sup>1</sup> Department of Pathobiology, College of Veterinary Medicine, University of Florida,  
Gainesville, Florida 32611-0880 and

<sup>2</sup> National Wildlife Health Center, 6006 Schroeder Road, Madison, Wisconsin 53711-6223  
(e-mail: rebecca\_cole@nbs.gov and constance\_roderick@nbs.gov)

**ABSTRACT:** A total of 28 species of helminths (17 trematodes, 3 cestodes, 7 nematodes, and 1 acanthocephalan) was recovered from 17 ospreys (*Pandion haliaetus*) from the United States. Intensities of infection were low and no lesions were attributed to the parasites. Seven species appear to be specialists in ospreys, 2 species generalists in raptors, and the remainder generalists in other orders of fish-eating birds. *Pandion-trema rjikovi*, *Diasiella diasi*, and *Contracaecum pandioni* are reported for the first time from North America.

**KEY WORDS:** helminths, osprey, parasites, *Pandion haliaetus*, *Pandion-trema rjikovi*, *Diasiella diasi*, *Contracaecum pandioni*.

The osprey, *Pandion haliaetus* (Linnaeus), is a cosmopolitan, monotypic member of the family Falconidae comprising its own subfamily, Pandioninae. Ospreys breed primarily in the Northern Hemisphere (North America and Eurasia) and winter in the Southern Hemisphere (South America, Africa, and India), with the exception of 2 nonmigratory subspecies in the Caribbean and Indonesia (Poole, 1989). Although this predominantly fish-eating raptor was considered threatened in North America in the 1960's because of pesticide contamination of the food chain, it has made a strong recovery and is now common in many parts of its former range (Ewins, 1994).

<sup>3</sup> Present address and author for reprints: 2108 Hilda Avenue, Missoula, Montana 59801.